This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problems Mailbox.

APPENDIX B

New York

Toronto

London

San Francisco

McGraw-Hill Book Company

UNIVERSITY OF CONNECTICUT

Leonid V. Azároff

PROFESSOR OF PHYSICS AND DIRECTOR

INSTITUTE OF MATERIALS SCIENCE

James J. Brophy

DIRECTOR OF TECHNICAL DEVELOPMENT IIT RESEARCH INSTITUTE processes

naterials

Metal Bond R&D

electronic

16/80/10 MED 12:48 LVX 1 208

700 🗗

PATENT

ELECTRONIC PROCESSES IN MATERIALS

Copyright © 1963 by McGraw-Hill, Inc. All Rights Reserved. Printed in the United States of America. This book, or parts thereof, may not be reproduced in any form without permission of the publishers. Library of Congress Catalog Card Number 62-21566

02669

6789101112 HDMM 7654321069

Control of the second of the s

Chapter 1

tance along the of a. Similarly, the length of b. es of all points oup. One com-Proceeding in a Fig. 22 are the ig b. Since it is coordinate along lext, the coordie $y\bar{x}$. The coorymmetry-related ated in Fig. 22. ry-related points an equipoint set. ing to this set, its

two points at ½0 the locations of 1 Fig. 22. Since s is "shared" by only two such he unit cell shown it is easy to see respectively, one. een therein, there re special positions et can take on any ontain the general

es of points

g, ÿx

:-dimensional space obtained by adding s shown in Table 4. structure of crystals


Since the symmetry of P4 does not affect z, this coordinate is quite general for all equipoints. Note that Table 4 contains an additional column headed Wyckoff notation. This notation was proposed in one of the first tabulations of such data, and it has become common practice to refer to a position in a unit cell by its rank and letter.

Table 4
Equipoints of space group P4

Rank of equipoint	Wyckoff notation	Symmetry of location	Coordinates of equivalent points
1 1 2 4	a. b c d	4 4 2 1	00z \frac{1}{2} \frac{1}{2}z 0\frac{1}{2}z, \frac{1}{2}0z xyz, y\frac{x}{2}z, \frac{x}{y}z, \tilde{y}xz

The closest packings of spheres

Hexagonal and cubic closest packings. To a very close approximation, the atoms in crystals can be likened to hard impenetrable spheres. It is instructive, therefore, to consider some of the ways that spheres can be packed together in three-dimensional arrays. As a start, consider the arrangement of like spheres in a plane. An array in which the maximum available space is occupied is shown in Fig. 23A. This array is called a






Fig. 23

hexagonal closest-packed layer, or simply a closest-packed layer, and it can be shown that the circles in Fig. 23A occupy 90.7 per cent of the available space (Exercise 10). The unit cell of the plane lattice of this array is shown by broken lines in Fig. 23A. Note that each unit cell contains two kinds of void spaces, labeled B and C in Fig. 23B. Note